

Section 4.5 Exponential and Logarithmic Equations

Exponential Equations

An *exponential equation* is one in which the variable occurs in the exponent.

EXAMPLE: Solve the equation $2^x = 7$.

Solution 1: We have

$$\begin{aligned}2^x &= 7 \\[\log_2 2^x &= \log_2 7] \\[x \log_2 2 &= \log_2 7] \\x &= \log_2 7 \approx 2.807\end{aligned}$$

Solution 2: We have

$$\begin{aligned}2^x &= 7 \\ \ln 2^x &= \ln 7 \\ x \ln 2 &= \ln 7 \\ x &= \frac{\ln 7}{\ln 2} \approx 2.807\end{aligned}$$

EXAMPLE: Solve the equation $4^{x+1} = 3$.

Solution 1: We have

$$\begin{aligned}4^{x+1} &= 3 \\[\log_4 4^{x+1} &= \log_4 3] \\[(x+1) \log_4 4 &= \log_4 3] \\x+1 &= \log_4 3 \\x &= \log_4 3 - 1 \approx -0.208\end{aligned}$$

Solution 2: We have

$$\begin{aligned}4^{x+1} &= 3 \\ \ln 4^{x+1} &= \ln 3 \\ (x+1) \ln 4 &= \ln 3 \\ x+1 &= \frac{\ln 3}{\ln 4} \\ x &= \frac{\ln 3}{\ln 4} - 1 \approx -0.208\end{aligned}$$

EXAMPLE: Solve the equation $3^{x-3} = 5$.

EXAMPLE: Solve the equation $3^{x-3} = 5$.

Solution 1: We have

$$3^{x-3} = 5$$

$$x - 3 = \log_3 5$$

$$x = \log_3 5 + 3 \approx 4.465$$

Solution 2: We have

$$3^{x-3} = 5$$

$$\ln 3^{x-3} = \ln 5$$

$$(x - 3) \ln 3 = \ln 5$$

$$x - 3 = \frac{\ln 5}{\ln 3}$$

$$x = \frac{\ln 5}{\ln 3} + 3 \approx 4.465$$

EXAMPLE: Solve the equation $8e^{2x} = 20$.

Solution: We have

$$8e^{2x} = 20$$

$$e^{2x} = \frac{20}{8} = \frac{5}{2}$$

$$2x = \ln \frac{5}{2}$$

$$x = \frac{\ln \frac{5}{2}}{2} = \frac{1}{2} \ln \frac{5}{2} \approx 0.458$$

EXAMPLE: Solve the equation $e^{3-2x} = 4$ algebraically and graphically.


Solution: We have

$$e^{3-2x} = 4$$

$$3 - 2x = \ln 4$$

$$-2x = \ln 4 - 3$$

$$x = \frac{\ln 4 - 3}{-2} = \frac{3 - \ln 4}{2} \approx 0.807$$



EXAMPLE: Solve the equation $e^{2x} - e^x - 6 = 0$.

Solution 1: We have

$$e^{2x} - e^x - 6 = 0$$

$$(e^x)^2 - e^x - 6 = 0$$

$$(e^x - 3)(e^x + 2) = 0$$

$$e^x - 3 = 0 \quad \text{or} \quad e^x + 2 = 0$$

$$e^x = 3$$

$$e^x = -2$$

The equation $e^x = 3$ leads to $x = \ln 3$. But the equation $e^x = -2$ has no solution because $e^x > 0$ for all x . Thus, $x = \ln 3 \approx 1.0986$ is the only solution.

Solution 1': Put $e^x = w$. Then

$$e^{2x} - e^x - 6 = 0$$

$$(e^x)^2 - e^x - 6 = 0$$

$$w^2 - w - 6 = 0$$

$$(w - 3)(w + 2) = 0$$

$$w - 3 = 0 \quad \text{or} \quad w + 2 = 0$$

$$w = 3$$

$$w = -2$$

$$e^x = 3$$

$$e^x = -2$$

The equation $e^x = 3$ leads to $x = \ln 3$. But the equation $e^x = -2$ has no solution because $e^x > 0$ for all x . Thus, $x = \ln 3 \approx 1.0986$ is the only solution.

EXAMPLE: Solve the equation $e^{2x} - 3e^x + 2 = 0$.

EXAMPLE: Solve the equation $e^{2x} - 3e^x + 2 = 0$.

Solution 1: We have

$$\begin{aligned}e^{2x} - 3e^x + 2 &= 0 \\(e^x)^2 - 3e^x + 2 &= 0 \\(e^x - 1)(e^x - 2) &= 0 \\e^x - 1 = 0 &\quad \text{or} \quad e^x - 2 = 0 \\e^x = 1 &\quad e^x = 2 \\x = 0 &\quad x = \ln 2\end{aligned}$$

Solution 1': Put $e^x = w$. Then

$$\begin{aligned}w^2 - 3w + 2 &= 0 \\(w - 1)(w - 2) &= 0 \\w - 1 = 0 &\quad \text{or} \quad w - 2 = 0 \\w = 1 &\quad w = 2 \\e^x = 1 &\quad e^x = 2 \\x = 0 &\quad x = \ln 2\end{aligned}$$

EXAMPLE: Solve the equation $7^{2x} - 3 \cdot 7^x + 1 = 0$.

Solution: Put $7^x = w$. Then

$$w^2 - 3w + 1 = 0 \implies w = \frac{-(-3) \pm \sqrt{(-3)^2 - 4 \cdot 1 \cdot 1}}{2 \cdot 1} = \frac{3 \pm \sqrt{5}}{2}$$

so $7^x = \frac{3 \pm \sqrt{5}}{2}$, therefore $x = \log_7 \left(\frac{3 \pm \sqrt{5}}{2} \right)$.

EXAMPLE: Solve the equation $7^{2x} - 7^x - 1 = 0$.

Solution: Put $7^x = w$. Then

$$w^2 - w - 1 = 0 \implies w = \frac{-(-1) \pm \sqrt{(-1)^2 - 4 \cdot 1 \cdot (-1)}}{2 \cdot 1} = \frac{1 \pm \sqrt{5}}{2}$$

Since $\frac{1 - \sqrt{5}}{2} < 0$, it follows that

$$7^x = \frac{1 + \sqrt{5}}{2} \implies x = \log_7 \left(\frac{1 + \sqrt{5}}{2} \right)$$

EXAMPLE: Solve the equation $3xe^x + x^2e^x = 0$.

Solution: We have

$$\begin{aligned}3xe^x + x^2e^x &= 0 \\xe^x(3 + x) &= 0 \\x(3 + x) &= 0 \\x = 0 &\quad \text{or} \quad 3 + x = 0 \\x = 0 &\quad x = -3\end{aligned}$$

Logarithmic Equations

A *logarithmic equation* is one in which a logarithm of the variable occurs.

EXAMPLE: Solve the equation $\ln x = 8$.

Solution: We have

$$\begin{aligned}\ln x &= 8 \\ [e^{\ln x} &= e^8] \\ x &= e^8\end{aligned}$$

EXAMPLE: Solve the equation $\log_2(x + 2) = 5$.

Solution: We have

$$\begin{aligned}\log_2(x + 2) &= 5 \\ [2^{\log_2(x+2)} &= 2^5] \\ x + 2 &= 2^5 \\ x &= 2^5 - 2 = 32 - 2 = 30\end{aligned}$$

EXAMPLE: Solve the equation $\log_7(25 - x) = 3$.

Solution: We have

$$\begin{aligned}\log_7(25 - x) &= 3 \\ [7^{\log_7(25-x)} &= 7^3] \\ 25 - x &= 7^3 \\ x &= 25 - 7^3 = 25 - 343 = -318\end{aligned}$$

EXAMPLE: Solve the equation $4 + 3\log(2x) = 16$.

EXAMPLE: Solve the equation $4 + 3 \log(2x) = 16$.

Solution: We have

$$4 + 3 \log(2x) = 16$$

$$3 \log(2x) = 12$$

$$\log(2x) = 4$$

$$2x = 10^4$$

$$x = \frac{10^4}{2} = \frac{10,000}{2} = 5,000$$

EXAMPLE: Solve the equation $\log(x + 2) + \log(x - 1) = 1$ algebraically and graphically.

Solution: We have

$$\log(x + 2) + \log(x - 1) = 1$$

$$\log[(x + 2)(x - 1)] = 1$$

$$(x + 2)(x - 1) = 10$$

$$x^2 + x - 2 = 10$$

$$x^2 + x - 12 = 0$$

$$(x + 4)(x - 3) = 0$$

$$x + 4 = 0$$

or

$$x - 3 = 0$$

$$x = -4$$


$$x = 3$$

We check these potential solutions in the original equation and find that $x = -4$ is not a solution (because logarithms of negative numbers are undefined), but $x = 3$ is a solution.

To solve the equation graphically we rewrite it as

$$\log(x + 2) + \log(x - 1) - 1 = 0$$

and then graph $y = \log(x + 2) + \log(x - 1) - 1$. The solutions are the x -intercepts of the graph.



EXAMPLE: Solve the following equations

(a) $\log(x + 8) + \log(x - 1) = 1$

(b) $\log(x^2 - 1) - \log(x + 1) = 3$

EXAMPLE: Solve the following equations

(a) $\log(x + 8) + \log(x - 1) = 1$

(b) $\log(x^2 - 1) - \log(x + 1) = 3$

Solution:

(a) We have

$$\log(x + 8) + \log(x - 1) = 1$$

$$\log[(x + 8)(x - 1)] = 1$$

$$(x + 8)(x - 1) = 10$$

$$x^2 + 7x - 8 = 10$$

$$x^2 + 7x - 18 = 0$$

$$(x + 9)(x - 2) = 0$$

$$x + 9 = 0$$

or

$$x - 2 = 0$$

$$x = -9$$

$$x = 2$$

We check these potential solutions in the original equation and find that $x = -9$ is not a solution (because logarithms of negative numbers are undefined), but $x = 2$ is a solution.

(b) We have

$$\log(x^2 - 1) - \log(x + 1) = 3$$

$$\log \frac{x^2 - 1}{x + 1} = 3$$

$$\frac{x^2 - 1}{x + 1} = 10^3$$


$$\frac{(x - 1)(x + 1)}{x + 1} = 1000$$

$$x - 1 = 1000$$

$$x = 1001$$

EXAMPLE: Solve the equation $x^2 = 2 \ln(x + 2)$ graphically.

Solution: We first move all terms to one side of the equation $x^2 - 2 \ln(x + 2) = 0$. Then we graph $y = x^2 - 2 \ln(x + 2)$. The solutions are the x -intercepts of the graph.



EXAMPLE: Find the solution of the equation, correct to two decimal places.

(a) $10^{x+3} = 6^{2x}$

(b) $5 \ln(3 - x) = 4$

(c) $\log_2(x + 2) + \log_2(x - 1) = 2$

EXAMPLE: Find the solution of the equation, correct to two decimal places.

(a) $10^{x+3} = 6^{2x}$

(b) $5 \ln(3 - x) = 4$

(c) $\log_2(x + 2) + \log_2(x - 1) = 2$

Solution:

(a) We have

$$10^{x+3} = 6^{2x}$$

$$\ln 10^{x+3} = \ln 6^{2x}$$

$$(x + 3) \ln 10 = 2x \ln 6$$

$$x \ln 10 + 3 \ln 10 = 2x \ln 6$$

$$x \ln 10 - 2x \ln 6 = -3 \ln 10$$

$$x(\ln 10 - 2 \ln 6) = -3 \ln 10$$

$$x = \frac{-3 \ln 10}{\ln 10 - 2 \ln 6} \approx 5.39$$

(b) We have

$$5 \ln(3 - x) = 4$$

$$\ln(3 - x) = \frac{4}{5}$$

$$3 - x = e^{4/5}$$

$$x = 3 - e^{4/5} \approx 0.77$$

(c) We have

$$\log_2(x + 2) + \log_2(x - 1) = 2$$

$$\log_2(x + 2)(x - 1) = 2$$

$$(x + 2)(x - 1) = 4$$

$$x^2 + x - 2 = 4$$

$$x^2 + x - 6 = 0$$

$$(x - 2)(x + 3) = 0$$

$$x - 2 = 0$$

or

$$x + 3 = 0$$

$$x = 2$$

$$x = -3$$

Since $x = -3$ is not from the domain of $\log_2(x + 2) + \log_2(x - 1)$, the only answer is $x = 2$.

Applications

EXAMPLE: If I_0 and I denote the intensity of light before and after going through a material and x is the distance (in feet) the light travels in the material, then according to the **Beer-Lambert Law**

$$-\frac{1}{k} \ln \left(\frac{I}{I_0} \right) = x$$

where k is a constant depending on the type of material.

(a) Solve the equation for I .

(b) For a certain lake $k = 0.025$ and the light intensity is $I_0 = 14$ lumens (lm). Find the light intensity at a depth of 20 ft.

Solution:

(a) We first isolate the logarithmic term.

$$-\frac{1}{k} \ln \left(\frac{I}{I_0} \right) = x$$

$$\ln \left(\frac{I}{I_0} \right) = -kx$$

$$\frac{I}{I_0} = e^{-kx}$$

$$I = I_0 e^{-kx}$$

(b) We find I using the formula from part (a).

$$I = I_0 e^{-kx} = 14e^{(-0.025)(20)} \approx 8.49$$

The light intensity at a depth of 20 ft is about 8.5 lm.

EXAMPLE: A sum of \$5000 is invested at an interest rate of 5% per year. Find the time required for the money to double if the interest is compounded according to the following method.

(a) Semiannual

(b) Continuous

EXAMPLE: A sum of \$5000 is invested at an interest rate of 5% per year. Find the time required for the money to double if the interest is compounded according to the following method.

- (a) Semiannual (b) Continuous

Solution:

- (a) We use the formula for compound interest

$$A(t) = P \left(1 + \frac{r}{n}\right)^{nt}$$

with $P = \$5000$, $A(t) = \$10,000$, $r = 0.05$, $n = 2$ and solve the resulting exponential equation for t .

$$5000 \left(1 + \frac{0.05}{2}\right)^{2t} = 10,000$$

$$(1.025)^{2t} = 2$$

$$\log 1.025^{2t} = \log 2$$

$$2t \log 1.025 = \log 2$$

$$t = \frac{\log 2}{2 \log 1.025} \approx 14.04$$

The money will double in 14.04 years.

- (b) We use the formula for continuously compounded interest

$$A(t) = Pe^{rt}$$

with $P = \$5000$, $A(t) = \$10,000$, $r = 0.05$ and solve the resulting exponential equation for t .

$$5000e^{0.05t} = 10,000$$

$$e^{0.05t} = 2$$

$$0.05t = \ln 2$$

$$t = \frac{\ln 2}{0.05} \approx 13.86$$

The money will double in 13.86 years.

EXAMPLE: A sum of \$1000 is invested at an interest rate of 4% per year. Find the time required for the amount to grow to \$4000 if interest is compounded continuously.

EXAMPLE: A sum of \$1000 is invested at an interest rate of 4% per year. Find the time required for the amount to grow to \$4000 if interest is compounded continuously.

Solution: We use the formula for continuously compounded interest

$$A(t) = Pe^{rt}$$

with $P = \$1000$, $A(t) = \$4000$, $r = 0.04$ and solve the resulting exponential equation for t .

$$1000e^{0.04t} = 4000$$

$$e^{0.04t} = 4$$

$$0.04t = \ln 4$$

$$t = \frac{\ln 4}{0.04} \approx 34.66$$

The amount will be \$4000 in about 34 years and 8 months.