

# The Fibonacci Numbers

DEFINITION: The **Fibonacci sequence** is defined recursively by  $f_1 = 1, f_2 = 1$ , and

$$f_n = f_{n-1} + f_{n-2} \quad \text{for } n \geq 3$$

The terms of this sequence are called the **Fibonacci numbers**.

The Fibonacci sequence begins with the integers

$$1, 1, 2, 3, 5, 8, 13, 21, 34, 55, 89, 144, \dots$$

REMARK: We can define the value  $f_0 = 0$  so that  $f_2 = f_1 + f_0$ .

EXAMPLE: For any positive integer  $n$ ,

$$\sum_{k=1}^n f_k = f_{n+2} - 1$$

Proof: Since

$$f_{k+2} = f_{k+1} + f_k$$

it follows that

$$f_k = f_{k+2} - f_{k+1}$$

therefore

$$\begin{aligned} \sum_{k=1}^n f_k &= \sum_{k=1}^n (f_{k+2} - f_{k+1}) \\ &= (f_3 - f_2) + (f_4 - f_3) + (f_5 - f_4) + \dots + (f_{n+1} - f_n) + (f_{n+2} - f_{n+1}) \\ &= f_{n+2} - f_2 \\ &= f_{n+2} - 1 \end{aligned}$$

EXAMPLE: For any positive integer  $n$ ,

$$f_{n+3} + f_n = 2f_{n+2} \tag{1}$$

Proof 1: We have

$$f_{n+3} + f_n = f_{n+2} + f_{n+1} + f_n = f_{n+2} + f_{n+2} = 2f_{n+2}$$

Proof 2: We have

$$f_n + f_{n+1} = f_{n+2} \tag{2}$$

and

$$f_{n+1} + f_{n+2} = f_{n+3} \tag{3}$$

Subtracting (3) from (2) we get

$$f_n - f_{n+2} = f_{n+2} - f_{n+3}$$

which can be rewritten as (1).

EXAMPLE: For any positive integer  $n$ ,

$$f_{2n} = f_{n+1}^2 - f_{n-1}^2 \quad (4)$$

Proof:

**STEP 1:** For  $n = 1$  (4) becomes

$$f_2 = f_2^2 - f_0^2$$

which is true, since  $1 = 1^2 - 0^2$ . Similarly, for  $n = 2$  (4) becomes

$$f_4 = f_3^2 - f_1^2$$

which is true, since  $3 = 2^2 - 1^2$ .

**STEP 2:** Suppose (4) is true for some  $n = k \geq 1$ , that is

$$f_{2k} = f_{k+1}^2 - f_{k-1}^2$$

and  $n = k + 1$ , that is

$$f_{2k+2} = f_{k+2}^2 - f_k^2$$

**STEP 3:** Prove that (4) is true for  $n = k + 2$ , that is

$$f_{2k+4} \stackrel{?}{=} f_{k+3}^2 - f_{k+1}^2$$

We have

$$\begin{aligned} f_{2k+4} &= f_{2k+3} + f_{2k+2} = f_{2k+2} + f_{2k+1} + f_{2k+2} = 2f_{2k+2} + f_{2k+1} \\ &= 2f_{2k+2} + f_{2k+2} - f_{2k} \\ &= 3f_{2k+2} - f_{2k} \\ &\stackrel{\text{ST.2}}{=} 3(f_{k+2}^2 - f_k^2) - (f_{k+1}^2 - f_{k-1}^2) \\ &\stackrel{?}{=} f_{k+3}^2 - f_{k+1}^2 \end{aligned}$$

We now show that which is true, since

$$\begin{aligned} &3(f_{k+2}^2 - f_k^2) - (f_{k+1}^2 - f_{k-1}^2) \stackrel{?}{=} f_{k+3}^2 - f_{k+1}^2 \\ &\quad \uparrow \\ &3(f_{k+2}^2 - f_k^2) - f_{k+1}^2 + f_{k-1}^2 \stackrel{?}{=} f_{k+3}^2 - f_{k+1}^2 \\ &\quad \uparrow \\ &3(f_{k+2}^2 - f_k^2) + f_{k-1}^2 \stackrel{?}{=} f_{k+3}^2 \\ &\quad \uparrow \\ &3(f_{k+2}^2 - f_k^2) \stackrel{?}{=} f_{k+3}^2 - f_{k-1}^2 \\ &\quad \uparrow \\ &3(f_{k+2} - f_k)(f_{k+2} + f_k) \stackrel{?}{=} (f_{k+3} - f_{k-1})(f_{k+3} + f_{k-1}) \\ &\quad \uparrow \\ &3f_{k+1}(f_{k+1} + f_k + f_k) \stackrel{?}{=} (f_{k+2} + f_{k+1} - f_{k-1})(f_{k+2} + f_{k+1} + f_{k-1}) \\ &\quad \uparrow \\ &3f_{k+1}(f_{k+1} + 2f_k) \stackrel{?}{=} (f_{k+1} + f_k + f_{k+1} - f_{k-1})(f_{k+1} + f_k + f_{k+1} + f_{k-1}) \\ &\quad \uparrow \\ &3f_{k+1}(f_{k+1} + 2f_k) = (f_{k+1} + f_k + f_k)(f_{k+1} + f_{k+1} + f_{k+1}) \end{aligned}$$

THEOREM 1: Let  $n$  be a positive integer  $\geq 3$  and let

$$\alpha = \frac{1 + \sqrt{5}}{2}$$

Then

$$f_n > \alpha^{n-2}$$

THEOREM 2: Let  $n$  be a positive integer and let

$$\alpha = \frac{1 + \sqrt{5}}{2} \quad \text{and} \quad \beta = \frac{1 - \sqrt{5}}{2}$$

Then

$$f_n = \frac{1}{\sqrt{5}}(\alpha^n - \beta^n)$$