

15

Multiple Integrals



Copyright © Cengage Learning. All rights reserved.

15.2

Iterated Integrals

Copyright © Cengage Learning. All rights reserved.

Iterated Integrals

Suppose that f is a function of two variables that is integrable on the rectangle $R = [a, b] \times [c, d]$.

We use the notation $\int_c^d f(x, y) dy$ to mean that x is held fixed and $f(x, y)$ is integrated with respect to y from $y = c$ to $y = d$. This procedure is called *partial integration with respect to y* . (Notice its similarity to partial differentiation.)

Now $\int_c^d f(x, y) dy$ is a number that depends on the value of x , so it defines a function of x :

$$A(x) = \int_c^d f(x, y) dy$$

3

Iterated Integrals

If we now integrate the function A with respect to x from $x = a$ to $x = b$, we get

$$\boxed{1} \quad \int_a^b A(x) dx = \int_a^b \left[\int_c^d f(x, y) dy \right] dx$$

The integral on the right side of Equation 1 is called an **iterated integral**. Usually the brackets are omitted. Thus

$$\boxed{2} \quad \int_a^b \int_c^d f(x, y) dy dx = \int_a^b \left[\int_c^d f(x, y) dy \right] dx$$

means that we first integrate with respect to y from c to d and then with respect to x from a to b .

4

Iterated Integrals

Similarly, the iterated integral

$$\boxed{3} \quad \int_c^d \int_a^b f(x, y) dx dy = \int_c^d \left[\int_a^b f(x, y) dx \right] dy$$

means that we first integrate with respect to x (holding y fixed) from $x = a$ to $x = b$ and then we integrate the resulting function of y with respect to y from $y = c$ to $y = d$.

Notice that in both Equations 2 and 3 we work *from the inside out*.

5

Example 1

Evaluate the iterated integrals.

$$(a) \int_0^3 \int_1^2 x^2 y dy dx \quad (b) \int_1^2 \int_0^3 x^2 y dx dy$$

Solution:

(a) Regarding x as a constant, we obtain

$$\begin{aligned} \int_1^2 x^2 y dy &= \left[x^2 \frac{y^2}{2} \right]_{y=1}^{y=2} \\ &= x^2 \left(\frac{2^2}{2} \right) - x^2 \left(\frac{1^2}{2} \right) \\ &= \frac{3}{2} x^2 \end{aligned}$$

6

Example 1 – Solution

cont'd

Thus the function A in the preceding discussion is given by $A(x) = \frac{3}{2}x^2$ in this example.

We now integrate this function of x from 0 to 3:

$$\begin{aligned}\int_0^3 \int_1^2 x^2 y \, dy \, dx &= \int_0^3 \left[\int_1^2 x^2 y \, dy \right] dx \\ &= \int_0^3 \frac{3}{2}x^2 \, dx \\ &= \left. \frac{x^3}{2} \right|_0^3 \\ &= \frac{27}{2}\end{aligned}$$

7

Example 1 – Solution

cont'd

(b) Here we first integrate with respect to x :

$$\begin{aligned}\int_1^2 \int_0^3 x^2 y \, dx \, dy &= \int_1^2 \left[\int_0^3 x^2 y \, dx \right] dy \\ &= \int_1^2 \left[\frac{x^3}{3} y \right]_{x=0}^{x=3} dy \\ &= \int_1^2 9y \, dy \\ &= 9 \left. \frac{y^2}{2} \right|_1^2 = \frac{27}{2}\end{aligned}$$

8

Iterated Integrals

Notice that in Example 1 we obtained the same answer whether we integrated with respect to y or x first.

In general, it turns out (see Theorem 4) that the two iterated integrals in Equations 2 and 3 are always equal; that is, the order of integration does not matter. (This is similar to Clairaut's Theorem on the equality of the mixed partial derivatives.)

9

Iterated Integrals

The following theorem gives a practical method for evaluating a double integral by expressing it as an iterated integral (in either order).

4 Fubini's Theorem If f is continuous on the rectangle $R = \{(x, y) \mid a \leq x \leq b, c \leq y \leq d\}$, then

$$\iint_R f(x, y) \, dA = \int_a^b \int_c^d f(x, y) \, dy \, dx = \int_c^d \int_a^b f(x, y) \, dx \, dy$$

More generally, this is true if we assume that f is bounded on R , f is discontinuous only on a finite number of smooth curves, and the iterated integrals exist.

10

Iterated Integrals

In the special case where $f(x, y)$ can be factored as the product of a function of x only and a function of y only, the double integral of f can be written in a particularly simple form.

To be specific, suppose that $f(x, y) = g(x)h(y)$ and $R = [a, b] \times [c, d]$.

Then Fubini's Theorem gives

$$\iint_R f(x, y) \, dA = \int_c^d \int_a^b g(x)h(y) \, dx \, dy = \int_c^d \left[\int_a^b g(x)h(y) \, dx \right] dy$$

11

Iterated Integrals

In the inner integral, y is a constant, so $h(y)$ is a constant and we can write

$$\int_a^b g(x)h(y) \, dx = \int_a^b h(y) \left(\int_a^b g(x) \, dx \right) = \int_a^b g(x) \, dx \int_c^d h(y) \, dy$$

since $\int_a^b g(x) \, dx$ is a constant.

Therefore, in this case, the double integral of f can be written as the product of two single integrals:

$$\iint_R g(x)h(y) \, dA = \int_a^b g(x) \, dx \int_c^d h(y) \, dy \quad \text{where } R = [a, b] \times [c, d]$$

12