

AREA AND VOLUME

THEOREM:

If A is a 2×2 matrix, the area of the parallelogram determined by the columns of A is $|\det A|$. If A is a 3×3 matrix, the volume of the parallelepiped determined by the columns of A is $|\det A|$.

CRAMER'S RULE

DEFINITION:

For any $n \times n$ matrix A and any \bar{b} in R^n , let $A_i(\bar{b})$ be the matrix obtained from A by replacing column i by the vector \bar{b} .

EXAMPLE:

Let $A = \begin{bmatrix} 2 & 1 & 3 \\ 1 & 0 & 4 \\ 0 & 0 & 5 \end{bmatrix}$, $\bar{b} = \begin{bmatrix} 3 \\ 8 \\ 9 \end{bmatrix}$. Then

$$A_1(\bar{b}) = \begin{bmatrix} 3 & 1 & 3 \\ 8 & 0 & 4 \\ 9 & 0 & 5 \end{bmatrix} \quad A_2(\bar{b}) = \begin{bmatrix} 2 & 3 & 3 \\ 1 & 8 & 4 \\ 0 & 9 & 5 \end{bmatrix}$$

$$A_3(\bar{b}) = \begin{bmatrix} 2 & 1 & 3 \\ 1 & 0 & 8 \\ 0 & 0 & 9 \end{bmatrix}$$

THEOREM (CRAMER'S RULE):

Let A be an invertible $n \times n$ matrix. For any \bar{b} in R^n , the unique solution \bar{x} of $A\bar{x} = \bar{b}$ has entries given by

$$x_i = \frac{\det A_i(\bar{b})}{\det A}, \quad i = 1, 2, \dots, n.$$

PROBLEM: Solve using Cramer's rule

$$\begin{cases} x_1 - 2x_2 = 1 \\ 3x_1 + 4x_2 = -7 \end{cases}$$

PROBLEM: Solve using Cramer's rule

$$\begin{cases} x_1 - 2x_2 = 1 \\ 3x_1 + 4x_2 = -7 \end{cases}$$

SOLUTION: We have

$$x_1 = \frac{\begin{vmatrix} 1 & -2 \\ -7 & 4 \end{vmatrix}}{\begin{vmatrix} 1 & -2 \\ 3 & 4 \end{vmatrix}} = \frac{4 - 14}{4 - (-6)} = \frac{-10}{10} = -1$$

$$x_2 = \frac{\begin{vmatrix} 1 & 1 \\ 3 & -7 \end{vmatrix}}{\begin{vmatrix} 1 & -2 \\ 3 & 4 \end{vmatrix}} = \frac{-7 - 3}{10} = \frac{-10}{10} = -1$$

FORMULA FOR A^{-1}

DEFINITION:

For any $n \times n$ matrix A , let A_{ij} be the submatrix of A , formed by deleting row i and column j .

EXAMPLE: Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 0 \end{bmatrix}$. Then

$$A_{11} = \begin{bmatrix} 5 & 6 \\ 8 & 0 \end{bmatrix} \quad A_{12} = \begin{bmatrix} 4 & 6 \\ 7 & 0 \end{bmatrix} \quad A_{13} = \begin{bmatrix} 4 & 5 \\ 7 & 8 \end{bmatrix}$$

$$A_{21} = \begin{bmatrix} 2 & 3 \\ 8 & 0 \end{bmatrix} \quad A_{22} = \begin{bmatrix} 1 & 3 \\ 7 & 0 \end{bmatrix} \quad A_{23} = \begin{bmatrix} 1 & 2 \\ 7 & 8 \end{bmatrix}$$

$$A_{31} = \begin{bmatrix} 2 & 3 \\ 5 & 6 \end{bmatrix} \quad A_{32} = \begin{bmatrix} 1 & 3 \\ 4 & 6 \end{bmatrix} \quad A_{33} = \begin{bmatrix} 1 & 2 \\ 4 & 5 \end{bmatrix}$$

THEOREM (AN INVERSE FORMULA):

Let A be an invertible $n \times n$ matrix.
Then

$$A^{-1} = \frac{1}{\det A} \begin{bmatrix} C_{11} & C_{12} & \dots & C_{1n} \\ C_{21} & C_{22} & \dots & C_{2n} \\ \dots & \dots & \dots & \dots \\ C_{n1} & C_{n2} & \dots & C_{nn} \end{bmatrix}^T,$$

where

$$C_{ij} = (-1)^{i+j} \det A_{ij}.$$

EXAMPLE: Let $A = \begin{bmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 0 \end{bmatrix}$. Find A^{-1} .

SOLUTION:

Step 1: One can verify that $\det A = 27$.

Step 2: We have

$$\begin{aligned} A_{11} &= \begin{bmatrix} 5 & 6 \\ 8 & 0 \end{bmatrix} & A_{12} &= \begin{bmatrix} 4 & 6 \\ 7 & 0 \end{bmatrix} & A_{13} &= \begin{bmatrix} 4 & 5 \\ 7 & 8 \end{bmatrix} \\ \det A_{11} &= -48 & \det A_{12} &= -42 & \det A_{13} &= -3 \\ C_{11} &= -48 & C_{12} &= 42 & C_{13} &= -3 \end{aligned}$$

$$\begin{aligned} A_{21} &= \begin{bmatrix} 2 & 3 \\ 8 & 0 \end{bmatrix} & A_{22} &= \begin{bmatrix} 1 & 3 \\ 7 & 0 \end{bmatrix} & A_{23} &= \begin{bmatrix} 1 & 2 \\ 7 & 8 \end{bmatrix} \\ \det A_{21} &= -24 & \det A_{22} &= -21 & \det A_{23} &= -6 \\ C_{21} &= 24 & C_{22} &= -21 & C_{23} &= 6 \end{aligned}$$

$$\begin{aligned} A_{31} &= \begin{bmatrix} 2 & 3 \\ 5 & 6 \end{bmatrix} & A_{32} &= \begin{bmatrix} 1 & 3 \\ 4 & 6 \end{bmatrix} & A_{33} &= \begin{bmatrix} 1 & 2 \\ 4 & 5 \end{bmatrix} \\ \det A_{31} &= -3 & \det A_{32} &= -6 & \det A_{33} &= -3 \\ C_{31} &= -3 & C_{32} &= 6 & C_{33} &= -3 \end{aligned}$$

Step 3:

$$\begin{aligned} A^{-1} &= \frac{1}{27} \begin{bmatrix} -48 & 42 & -3 \\ 24 & -21 & 6 \\ -3 & 6 & -3 \end{bmatrix}^T = \frac{1}{27} \begin{bmatrix} -48 & 24 & -3 \\ 42 & -21 & 6 \\ -3 & 6 & -3 \end{bmatrix} \\ &= \begin{bmatrix} -16/9 & 8/9 & -1/9 \\ 14/9 & -7/9 & 2/9 \\ -1/9 & 2/9 & -1/9 \end{bmatrix} \end{aligned}$$