

VECTOR SPACES:

$$1. \mathbf{R}^n = \left\{ \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} : x_1, \dots, x_n \in \mathbf{R} \right\}$$

2. The set P_n of polynomials of degree at most n :

$$\bar{p}(t) = a_n t^n + \dots + a_2 t^2 + a_1 t + a_0$$

where the coefficients a_n, \dots, a_0 and the variable t are real numbers.

3. The set of all real-valued functions defined on \mathbf{R} .

STANDARD BASIS FOR R^n :

$$\bar{e}_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad \bar{e}_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \dots, \quad \bar{e}_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}$$

STANDARD BASIS FOR P_n :

Vectors

$\bar{e}_1 = 1, \bar{e}_2 = t, \bar{e}_3 = t^2, \dots, \bar{e}_{n+1} = t^n$
form the so-called standard basis for the
vector space P_n .

DEFINITION:

Suppose $B = \{\bar{b}_1, \dots, \bar{b}_n\}$ is a basis for a vector space V and \bar{x} is in V . The coordinates of \bar{x} relative to the basis B are the weights c_1, \dots, c_n such that

$$\bar{x} = c_1\bar{b}_1 + \dots + c_n\bar{b}_n.$$

NOTATION:

$$[\bar{x}]_B = \begin{bmatrix} c_1 \\ \dots \\ c_n \end{bmatrix}$$

PROBLEM:

Let $B = \{1, t, t^2\}$ be the standard basis for P_2 . Find coordinates of the vector

$$\bar{p}(t) = -4 + 3t - 5t^2$$

relative to B .

SOLUTION:

By the definition above we have:

$$[\bar{p}]_B = \begin{bmatrix} -4 \\ 3 \\ -5 \end{bmatrix}.$$

PROBLEM:

Determine whether the polynomial

$$\bar{p}(t) = 2 + t + 7t^2 + 5t^3$$

can be represented as a linear combination of the polynomials

$$1 + t + 4t^2 + 3t^3, \quad 2 - t + 5t^2 + 3t^3.$$

SOLUTION:

The answer is “Yes”, because

$$\begin{bmatrix} 1 & 2 & 2 \\ 1 & -1 & 1 \\ 4 & 5 & 7 \\ 3 & 3 & 5 \end{bmatrix} \sim \begin{bmatrix} 1 & 2 & 2 \\ 0 & 3 & 1 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix},$$

and the echelon form of the augmented matrix represents a consistent system.

PROBLEM 1:

Determine whether the polynomials

$$1 + t^3, \quad 3 + t - 2t^2, \quad -t + 3t^2 - t^3$$

are linearly independent.

PROBLEM 2:

Determine whether the polynomials

$$1 - 3t + 5t^2, \quad -3 + 5t - 7t^2, \quad -4 + 5t - 6t^2, \quad 1 - t^2$$

span P_2 .

PROBLEM 3:

Determine whether the polynomials

$$3 + 7t, \quad 5 + t - 2t^3, \quad t - 2t^2, \quad 1 + 16t - 6t^2 + 2t^3$$

form a basis for P_3 .

PROBLEM 4:

Determine whether the polynomials

$$1 + t, \quad 1 + t^2, \quad t + t^2$$

form a basis for P_2 . If “Yes”, find coordinates of the vector

$$\bar{p}(t) = -4 + 3t - 5t^2$$

relative to this basis.

Solution of Problem 1:

Let $B = \{1, t, t^2, t^3\}$ be the standard basis of P_3 . Then polynomials

$$1 + t^3, \quad 3 + t - 2t^2, \quad -t + 3t^2 - t^3$$

produce coordinate vectors

$$\begin{bmatrix} 1 \\ 0 \\ 0 \\ 1 \end{bmatrix}, \quad \begin{bmatrix} 3 \\ 1 \\ -2 \\ 0 \end{bmatrix}, \quad \begin{bmatrix} 0 \\ -1 \\ 3 \\ -1 \end{bmatrix}$$

relative to B . Writing these vectors as the columns of a matrix A , we can determine their independence:

$$\begin{bmatrix} 1 & 3 & 0 \\ 0 & 1 & -1 \\ 0 & -2 & 3 \\ 1 & 0 & -1 \end{bmatrix} \sim \begin{bmatrix} 1 & 3 & 0 \\ 0 & 1 & -1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix}.$$

Since there are pivots in every column,

$$1 + t^3, \quad 3 + t - 2t^2, \quad -t + 3t^2 - t^3$$

are linearly independent.

Solution of Problem 2:

Let $B = \{1, t, t^2\}$ be the standard basis of P_2 . Then polynomials

$1-3t+5t^2$, $-3+5t-7t^2$, $-4+5t-6t^2$, $1-t^2$

produce coordinate vectors

$$\begin{bmatrix} 1 \\ -3 \\ 5 \end{bmatrix}, \begin{bmatrix} -3 \\ 5 \\ -7 \end{bmatrix}, \begin{bmatrix} -4 \\ 5 \\ -6 \end{bmatrix}, \begin{bmatrix} 1 \\ 0 \\ -1 \end{bmatrix}$$

relative to B . We have:

$$\begin{bmatrix} 1 & -3 & -4 & 1 \\ -3 & 5 & 5 & 0 \\ 5 & -7 & -6 & -1 \end{bmatrix} \sim \begin{bmatrix} 1 & -3 & -4 & 1 \\ 0 & 4 & 7 & -3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since there are 2 pivots and 3 rows, the polynomials

$1-3t+5t^2$, $-3+5t-7t^2$, $-4+5t-6t^2$, $1-t^2$

do not span P_2 .

Solution of Problem 3:

Let $B = \{1, t, t^2, t^3\}$ be the standard basis of P_3 . Then polynomials

$$3+7t, 5+t-2t^3, t-2t^2, 1+16t-6t^2+2t^3$$

produce coordinate vectors

$$\begin{bmatrix} 3 \\ 7 \\ 0 \\ 0 \end{bmatrix}, \begin{bmatrix} 5 \\ 1 \\ 0 \\ -2 \end{bmatrix}, \begin{bmatrix} 0 \\ 1 \\ -2 \\ 0 \end{bmatrix}, \begin{bmatrix} 1 \\ 16 \\ -6 \\ 2 \end{bmatrix}$$

relative to B . We have:

$$\begin{bmatrix} 3 & 5 & 0 & 1 \\ 7 & 1 & 1 & 16 \\ 0 & 0 & -2 & -6 \\ 0 & -2 & 0 & 2 \end{bmatrix} \sim \begin{bmatrix} 3 & 5 & 0 & 1 \\ 0 & 32 & -3 & -41 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Since there are 3 pivots and 4 columns, the polynomials

$$3+7t, 5+t-2t^3, t-2t^2, 1+16t-6t^2+2t^3$$

do not form a basis for P_3 .

Solution of Problem 4:

Let $B = \{1, t, t^2\}$ be the standard basis of P_2 . Then polynomials

$$1 + t, 1 + t^2, t + t^2$$

produce coordinate vectors

$$\begin{bmatrix} 1 \\ 1 \\ 0 \end{bmatrix}, \quad \begin{bmatrix} 1 \\ 0 \\ 1 \end{bmatrix}, \quad \begin{bmatrix} 0 \\ 1 \\ 1 \end{bmatrix}$$

relative to B . We have:

$$\begin{bmatrix} 1 & 1 & 0 \\ 1 & 0 & 0 \\ 0 & 1 & 1 \end{bmatrix} \sim \begin{bmatrix} 1 & 1 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}.$$

Since there are 3 pivots and 3 columns, the polynomials

$$1 + t, 1 + t^2, t + t^2$$

form a basis for P_2 .

Let

$$B = \{1 + t, 1 + t^2, t + t^2\}.$$

To find coordinates of the vector

$$\bar{p}(t) = -4 + 3t - 5t^2$$

relative to B , we consider the augmented matrix

$$\begin{bmatrix} 1 & 1 & 0 & -4 \\ 1 & 0 & 1 & 3 \\ 0 & 1 & 1 & -5 \end{bmatrix} \sim \begin{bmatrix} 1 & 0 & 0 & 2 \\ 0 & 1 & 0 & -6 \\ 0 & 0 & 1 & 1 \end{bmatrix},$$

therefore

$$[\bar{p}]_B = \begin{bmatrix} 2 \\ -6 \\ 1 \end{bmatrix}.$$